

Gila County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Apache (Arizona) trout	<i>Oncorhynchus apache</i>	Threatened	This yellowish or yellow-olive cutthroat-like trout has large dark spots on body. Its dorsal, anal, and caudal fins are edged with white. It has no red lateral band.	Apache, Coconino, Gila, Graham, Greenlee, Navajo	>5000 ft	Presently restricted to cold mountain streams with many low gradient meadow reaches.	Occupies stream habitats with substrates of boulders, rocks, and gravel with some sand or silt through mixed conifer and spruce-fir forests, and montane meadows and grasslands in the White Mountains. Also managed as a sport fish under special regulations.
Arizona hedgehog	<i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i>	Endangered	Dark green cylindroid 2.5-12 inches tall, 2-10 inches in diameter, single or in clusters. 1-3 gray or pinkish central spines largest deflexed and 5-11 shorter radial spines. Flower: brilliant red, side of stem in April-May.	Gila, Pinal	3,700-5,200 ft	Ecotone between interior chaparral and madrean evergreen woodland.	Open slopes, in narrow cracks between boulders, and in understory of shrubs. Additional genetic studies have determined that the species does not occur outside of the type locality.
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large, adults have white head and tail. Height 28-38 inches; wingspan 66-96 inches. Dark with varying degrees of mottled brown plumage. Feet bare of feathers.	Apache, Cochise, Coconino, Gila, Graham, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	Varies	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Some birds are nesting residents while a larger number winters along rivers and reservoirs. An estimated 200 to 300 birds winter in Arizona. Once endangered (32 FR 4001, 03-11-1967; 43 FR 6233, 02-14-78) because of reproductive failures from pesticide poisoning and loss of habitat, this species was down listed to threatened on August 11, 1995. Illegal shooting, disturbance, and loss of habitat continues to be a problem. Species has been proposed for delisting (64 FR 36454) but still receives full protection under the ESA.
California Brown pelican	<i>Pelecanus occidentalis californicus</i>	Endangered	Large dark gray-brown water bird with a pouch underneath long bill and webbed feet. Adults have a white head and neck, brownish black breast, and silver gray upper parts.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	Varies	Coastal land and islands; species found around many Arizona lakes and rivers.	Subspecies is found on Pacific Coast and is endangered due to pesticides. It is an uncommon transient in Arizona on many Arizona lakes and rivers. Individuals wander up from Mexico in summer and fall. No breeding records in Arizona.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Chiricahua leopard frog	<i>Rana chiricahuensis</i>	Threatened	Cream colored tubercles (spots) on a dark background on the rear of the thigh, dorsolateral folds that are interrupted and deflected medially, and a call given out of water distinguish this spotted frog from other leopard frogs.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Navajo, Pima, Santa Cruz, Yavapai	3300-8900 ft	Streams, rivers, backwaters, ponds, and stock tanks that are mostly free from introduced fish, crayfish, and bullfrogs.	Require permanent or nearly permanent water sources. Populations north of the Gila River may be a closely-related, but distinct, undescribed species. A special rule allows take of frogs due to operation and maintenance of livestock tanks on State and private lands.
Colorado pikeminnow	<i>Ptychocheilus lucius</i>	Endangered	Largest american minnow (up to 6 feet and 80 lbs) dusky-green, slender body has gold flecks on the dorsal surface. Head long and slender.	Gila, Yavapai	< 4,000 ft	Warm, swift, turbid mainstem rivers. Prefers eddies and pools.	Experimental non-essential (treated as proposed threatened). No critical habitat in Arizona.
Gila chub	<i>Gila intermedia</i>	Endangered	Deep compressed body, flat head. Dark olive-gray color above, silver sides. Endemic to Gila River Basin.	Cochise, Gila, Graham, Greenlee, Maricopa, Pima, Pinal, Santa Cruz, Yavapai	2,000 - 5,500 ft	Pools, springs, cienegas, and streams.	Found on multiple private lands, including the Nature Conservancy, the Audubon Society, and others. Also occurs on Federal and state lands and in Sonora, Mexico. Critical habitat occurs in Cochise, Gila, Graham, Greenlee, Pima, Pinal, Santa Cruz and Yavapai counties.
Gila topminnow	<i>Poeciliopsis occidentalis occidentalis</i>	Endangered	Small (2 inches), guppy-like, live bearing, lacks dark spots on its fins. Breeding males are jet black with yellow fins.	Gila, Graham, La Paz, Maricopa, Pima, Pinal, Santa Cruz, Yavapai	< 4,500 ft	Small streams, springs, and cienegas vegetated shallows.	Species historically occurred in backwaters of large rivers but is currently isolated to small streams and springs.
Gila trout	<i>Oncorhynchus gilae</i>	Threatened	Deep bodied with fine profuse spotting on the body, dorsal, and adipose fins. Dorsal, anal, and pelvic fins edged in white.	Gila, Greenlee	5,000-10,000 ft	Small high mountain streams.	Fish stocked into Dude Creek in Sept 1999 and into Raspberry Creek in Nov 2000. Also occurs in New Mexico.
Lesser long-nosed bat	<i>Leptonycteris curasoae yerbabuenae</i>	Endangered	Elongated muzzle, small leaf nose, and long tongue. Yellowish brown or gray above and cinnamon brown below. Tail minute and appears to be lacking. Easily disturbed.	Cochise, Gila, Graham, Greenlee, Pima, Pinal, Maricopa, Santa Cruz	< 6000 ft	Desert scrub habitat with agave and columnar cacti present as food plants.	Day roosts in caves and abandoned tunnels. Forages at night on nectar, pollen, and fruit of paniculate agaves and columnar cacti. This species is migratory and is present in Arizona usually from April to September and south of the border the remainder of the year.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Loach minnow	<i>Tiaroga cobitis</i>	Threatened	Small (<3 inches) slender, elongated fish, olive colored with dirty white spots at the base of the dorsal and caudal fins. Breeding males vivid red on mouth and base of fins.	Apache, Graham, Greenlee, Pinal, Navajo, Gila	<8000 ft	Benthic species of small to large perennial streams with swift shallow water over cobble and gravel. Recurrent flooding and natural hydrograph important.	Presently found in Aravaipa Creek, Deer Creek, Turkey Creek, Blue River, Campbell Blue Creek, San Francisco River, Eagle Creek, North Fork East Fork Black River, and White River in Arizona, and Dry Blue Creek, Pace Creek, Frieborn Creek, the Tularosa River, West Fork Gila River, and the mainstem upper Gila River in New Mexico. Proposed critical habitat (70 FR 75545-75590, December 20, 2005) includes portions of East Fork Black River, North Fork East Fork Black River, Boneyard Creek, Aravaipa Creek, Turkey Creek, Deer Creek, Eagle Creek, San Francisco River, Blue River, Campbell Blue Creek, and Little Blue Creek found in Apache, Graham, Greenlee, and Pinal counties, Arizona, as well as portions of the Blue River, San Francisco River, Tularosa River, Negrito Creek, Pace Creek, Dry Blue Creek, Frieborn Creek, Whitewater Creek, Gila River, and its West, Middle, and East Forks in Catron, Grant, and Hidalgo counties in New Mexico.
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Medium sized with dark eyes and no ear tufts. Brownish and heavily spotted with white or beige.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai	4100-9000 ft	Nests in canyons and dense forests with multi-layered foliage structure.	Generally nest in older forests of mixed conifer or ponderosa pine/gambel oak type, in canyons, and use variety of habitats for foraging. Sites with cool microclimates appear to be of importance or are preferred. Critical habitat was finalized on August 31, 2004 (69 FR 53182). Critical habitat in Arizona occurs in Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Navajo, Pima, Pinal, Santa Cruz, and Yavapai counties.
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Species is also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam; Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizona/New Mexico border to Coolidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	<8500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886) and can be viewed at http://arizonaes.fws.gov . In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.
Spikedace	<i>Meda fulgida</i>	Threatened	Small (<3 inches) slim with silvery sides and "spine" on dorsal fin. Breeding males brassy golden color.	Graham, Greenlee, Gila, Navajo, Pinal, Yavapai	< 6000 ft	Moderate to large perennial streams with gravel cobble substrates and moderate to swift velocities over sand and gravel substrates. Recurrent flooding and natural hydrograph important.	Presently found in Aravaipa Creek, Eagle Creek, Verde River, and the Gila River form the San Pedro River to Ashurst-Hayden Dam in Arizona, and the Gila River and its East and West Forks in New Mexico. Proposed critical habitat (70 FR 75545-75590, December 20, 2005) includes portions of the Verde River, Gila River, lower San Pedro River, Aravaipa Creek, and Eagle Creek in Graham, Greenlee, Pinal, and Yavapai counties in Arizona, and the Gila River and its East, Middle, and West Forks in Catron, Grant, and Hidalgo counties in New Mexico.
Yuma clapper rail	<i>Rallus longirostris yumanensis</i>	Endangered	Water bird with long legs and short tail. Long, slender decurved bill. Mottled brown or gray on its rump. Flanks and undersides are dark gray with narrow vertical stripes producing a barring effect.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	< 4,500 ft	Fresh water and brackish marshes.	Species is associated with dense emergent riparian vegetation. Requires wet substrate (mudflat, sandbar) with dense herbaceous or woody vegetation for nesting and foraging. Channelization and marsh destruction are primary sources of habitat loss.
Headwater chub	<i>Gila nigra</i>	Candidate	A streamlined dark gray to brown fish, often with longitudinal stripes on the sides, reaching a maximum size of about 12 inches.	Yavapai, Gila, Graham	3,000 - 6,700 ft	Small to medium-sized streams, often associated with deep pools and cover such as boulders or vegetation.	Occurs in the East Verde River and tributaries, Fossil Creek, Wet Bottom Creek, Deadman Creek, Tonto Creek and tributaries, San Carlos River, Ash Creek, and the upper Gila River in New Mexico.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill, which is blue-black with yellow on the lower half of the bill. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Listing was found warranted, but precluded as a distinct vertebrate population segment in the western U.S. on July 25, 2001. This finding indicates that the Service has sufficient information to list the bird, but other, higher priority listing actions prevent the Service from addressing the listing of the cuckoo at this time.
Arizona bugbane	<i>Cimicifuga arizonica</i>	Conservation Agreement	Perennial herb in the buttercup family up to 6-7 feet tall. Small white petal-less flowers appear in July-August. Fruit a follicle that splits open on one side as it dries.	Coconino, Gila	5,300-7,000 ft	Moist, loamy soil between coniferous and riparian ecotones.	Rich, fertile soils high in humus content, deep shade, and high humidity appears to be primary habitat requirements for this species. Conservation Agreement signed in June 1999.